


Monodisperse HPLC Particles




The Evolution of HPLC Columns

EVOSPHERE





EVO SPHERE



Monodisperse HPLC Columns

Fortis Technologies has designed a new fully porous monodisperse particle for use in HPLC columns. Combining this with a new range of selectivities gives the analyst the ability for high resolution, high efficiency separations.

Based upon a fully porous silica monodisperse particle, Evosphere® is the evolution of particle technology.

Combine a high efficiency particle with low backpressure, high loadability, scaleability and reproducibility and you have the ultimate combination.

Then add in novel selectivity options to provide enhanced resolution and selectivity and you have the capability to separate

more compounds in less time with greater sensitivity.

By building on a pure silica substrate method development and method transfer become more robust and reproducible across platforms as you scale from capillary to preparative.



Monodisperse Particles




Particle Morphology


Evosphere silica particles are manufactured to provide a high degree of monodispersity with a uniform smooth surface. Monodispersity generates high efficiency HPLC columns due to the reduced flow path dispersion (Eddy diffusion)

SEM imagery of the Evosphere in comparison with traditional particles highlights the much narrower size distribution.

Monodisperse Evosphere particles are available in 1.7 μm , 3 μm and 5 μm particle sizes.



Traditional
porous particles



Monodisperse
porous particles


Particle size distribution (D90/10)

When assigning a measurement to characterise a particle size distribution the ratio of D90/10 is often quoted, and as such can be used to gauge the degree of size uniformity of the particles.

The parameter D90 signifies the point in the size distribution, up to and including which, 90% of the total volume of material in the sample is 'contained'. For example; if the D90 is 6 μm , this means that 90% of the sample has size of 6 μm or smaller. The definition for D50, is then the size point below which 50% of the material is contained. Similarly, the D10 is the size below which 10% of the material is contained. This description has long been used in size distribution measurements.

As the particle size distribution for chromatographic silica moves towards monodisperse then the D90 and D10 values become closer together and the D90/10 value tends towards a value of 1.

Particle Size Distribution



	Monodisperse Silica	Commercial 3 μm Silica-A	Commercial 3 μm Silica-B
Median Particle size (d ₅₀)*	2.66 μm *	2.48 μm	2.97 μm
SEM Particle Size	3.0 μm	2.8 μm	3.3 μm
D90/10	1.12	1.58	1.61
Pore Volume	0.89	0.88	0.89

* Measured by Coulter Counter

Monodisperse Particles




Efficiency of Monodisperse particles

Analysts have had two ways of improving efficiency in the past. Move to a smaller particle with associated high backpressure and the need to buy a UHPLC instrument, or move to core-shell particles but with a compromise in loading and scalability.

Evosphere fully porous monodisperse particles have vastly increased efficiency over equivalent porous particle sizes. Due to maintaining high surface area, loading and retention time are not compromised as seen with core-shell particles.

- High Efficiency
- High Loading
- Scalable - capillary to Prep
- Robust
- Reproducible



Typical Efficiencies of HPLC particles




Efficiency Gains

The monodisperse nature of Evosphere coupled with our ability to pack the column more efficiently allows for a significant increase in efficiency and therefore resolution over traditional silica particles.


3µ Monodisperse vs 3µ Traditional



Efficiency



Loading Capacity





Fortis Evosphere® has a high surface area ($350\text{m}^2/\text{g}$) as per many modern Type B porous silica's, this allows loadability of compounds to be increased for purification purposes.

If you compare this to core-shell particles which typically have a surface area in the region of $130\text{m}^2/\text{g}$ you will quickly see overload and compromised peak shapes, meaning scale up of methods can be difficult.

Evosphere is available from capillary scale dimensions all the way up to preparative columns.


Comparison of loading capacity



Hydrophobicity


Evosphere® being a fully porous silica particle maintains a high surface area ($350\text{m}^2/\text{g}$) this means high retentive effect for analytes, as well as high loading capacity.

Evosphere C12 shows high hydrophobicity due to its high surface coverage, made possible by less steric hindrance when bonded.



Backpressure

Column impedance highlights how backpressure will remain the same for monodisperse particles as it will for polydisperse traditional particles. Backpressure in identically packed columns will only be effected by the nominal particle size.




Stationary Phase Choice




- Evosphere C18/PFP
- Orthogonal Selectivity
- Method development starting point

Evosphere C18/PFP is designed to provide characteristics which will enhance method development. It provides the ability to obtain sharp peak shapes whilst retaining and separating a wide variety of compounds both hydrophobic and hydrophilic.



- Evosphere C18/AR
- Orthogonal Selectivity
- Method development starting point

Evosphere C18/AR is designed to provide increased resolution between compounds, having a combination of hydrophobicity and aromatic selectivity will lead to enhanced resolution. USP L1 column.



- Evosphere RP18-Amide
- Orthogonal Selectivity
- Excellent method development option

Evosphere RP18-Amide is designed to provide polar characteristics which will enhance resolution in method development. It provides orthogonal selectivity to alkyl chain phases due to its polar-embedded group. Sharp peak shapes, extra selectivity and increased retention can all be obtained.




- Evosphere Diphenyl
- Separate positional isomers
- Stable ligand, No “MS” bleed

Evosphere Diphenyl is designed to provide pi-pi, steric and hydrophobic characteristics which will enhance selectivity and the ability to develop methods. Particularly suited to positional isomers and other closely related species such as metabolites.



- Evosphere Phenyl-Hexyl
- Separate metabolites
- Excellent resolution

Evosphere Phenyl-Hexyl is designed to provide characteristics which will enhance selectivity. It provides alternate selectivity to a pure hydrophobic stationary phase whilst still maintaining the key attributes of robustness and reproducibility.



- Evosphere AQUA
- Separate polar species
- Excellent stability

Evosphere AQUA is designed to provide characteristics which will enhance retention of highly polar analytes. Reproducible surface characteristics provide robust separations. Combination of hydrophobic and hydrophilic nature.



- Evosphere C12
- Ultra High Efficiency
- Method development starting point

Evosphere C12 is designed to provide characteristics which will enhance method development. The dense C12 ligand provides the ability to obtain sharp peak shapes whilst retaining and separating a wide variety of acid, base and neutral compounds with excellent robustness.



- Evosphere PFP
- Orthogonal Selectivity
- Combined with Ultra High Efficiency particles

Evosphere PFP (PentaFluoroPhenyl) is designed to provide characteristics which will enhance selectivity. It provides alternate selectivity to a hydrophobic stationary phase whilst still maintaining the key attributes of robustness and reproducibility.



- Evosphere HILIC
- High polarity compounds
- Combined with Ultra High Efficiency particles

Evosphere HILIC is designed to provide characteristics which will enhance retention of polar compounds. Hydrophilic interaction chromatography uses partition, ion-exchange and hydrogen bonding to retain high polarity analytes.

Evosphere Monodisperse Particles

	Particle Size	Surface Area	Pore Size	% C	pH range	USP
Evosphere C12	1.7µm 3µm 5µm	350m ² /g	100Å	17	1-11	L87
Evosphere C18/AR	1.7µm 3µm 5µm	350m ² /g	100Å	17	2-9	L1
Evosphere C18/PFP	1.7µm 3µm 5µm	350m ² /g	100Å	17	2-9	L1
Evosphere RP18-Amide	1.7µm 3µm 5µm	350m ² /g	100Å	20	2-9	L60
Evosphere Phenyl-Hexyl	1.7µm 3µm 5µm	350m ² /g	100Å	14	2-9	L11
Evosphere Diphenyl	1.7µm 3µm 5µm	350m ² /g	100Å	15	2-9	L11
Evosphere PFP	1.7µm 3µm 5µm	350m ² /g	100Å	13	2-9	L43
Evosphere AQUA	1.7µm 3µm 5µm	350m ² /g	100Å	18	2-9	L96
Evosphere HILIC	1.7µm 3µm 5µm	350m ² /g	100Å	n/a	2-7	L3

Principle Component Analysis (PCA)




PCA analysis

Principle component analysis has been used to differentiate columns for use in Chromatography for a number of years,* first developed by Euerby and Petersson based around tests suggested by Tanaka. A systematic approach to column characterisation allows the analyst to choose from a diverse (or similar) range of columns.

* M.Euerby, P.Petersson, LC-GC Europe (Sept 2000) 665-677

PCA analysis has been shown below to highlight the diversity of the stationary phases in the EvoSphere range.
Choose a phase based on orthogonal selectivity to your current column or by mechanism to match the analytes




EvoSphere PCA

Factors characterised:

- **k' PB** Retention factor of Pentylbenzene, an indicator of ligand density
- **Steric selectivity** Retention factor ratio between triphenylene and o-terphenyl $\alpha_{T/O} = k_1/k_0$ this is a measure of the shape selectivity and functionality of the silylating reagent.
- **Hydrophobic selectivity** α_{CH_2} Retention factor between pentylbenzene and butylbenzene. A measure of the surface coverage of the phase, differentiated by one methylene group is dependant upon the ligand density.
- **Hydrogen bonding capacity** $\alpha_{C/P}$ retention factor between caffeine and phenol. A indicator of the degree of endcapping
- **Ion-exchange capacity** α_{k_p/k_o} Retention factor between benzylamine and phenol at both pH 2.7 (acidic activity of silanols) and pH 7.6 an estimate of total silanol activity.

In the example below PCA analysis can be used to show the differences between several commercial Phenyl type phases. It can be seen how the new C18-PFP phase adds a new dimension being predominantly hydrophobic in nature but offering alternate selectivity due to the mixed C18 and PFP ligands.

EvoSphere Phenyl-Hex also offers a new selectivity to the range combining hydrophobicity and pi-pi interactions together.



HPLC/UHLC Phase Selectivity Chart

POLAR

EVOSPHERE HILIC

Hydrophilic Interaction chromatography

Retention of very high polarity analytes by partition, ion-exchange and hydrogen bonding

EVOSPHERE AQUA

Polar
Endcapped C18

Amino acids
Organic acids
High aqueous mobile phases

EVOSPHERE PFP

Pentafluorophenyl
Selectivity

Selectivity of aromatic compounds.
Orthogonal fluorine enhanced

EVOSPHERE RP18-AMIDE

Polar
Embedded C18

Strong acid retention
Sharp peak shape for basics

EVOSPHERE DIPHENYL

Dual Phenyl for
Steric selectivity

Isomers
Metabolites
Closely related species

EVOSPHERE PHENYL-HEXYL

Balanced aromatic and
hydrophobic selectivity

Balanced hydrophobic and
aromatic selectivity.
Hydrogen accepting for acids

EVOSPHERE C18/PFP

Pentafluorophenyl
and C18 ligands

Enhanced retention with
electron-rich molecules

EVOSPHERE C18/AR

Balanced Phenyl
and C18 ligands

Aromatic as well as hydrophobic
retention

HYDROPHOBIC


EVOSPHERE C12

C12 high coverage
stable ligand

Most Hydrophobic
Sharper peak shapes
High pH stable

Evosphere Selectivity

When developing a new method in chromatography having a diverse range of selectivities to choose from can help in deciding how peaks resolve and which is the best starting point. In this example a gradient run across several stationary phase shows orthogonal selectivity for many of the peaks.



1. Hydroquinone
2. Theobromine
3. Paracetamol
4. Theophylline
5. Paraxanthine
6. 4-Hydroxybenzoic acid
7. 2-Acetamidophenol
8. Caffeine
9. Phenol
10. Aspirin
11. 2-hydroxybenzoic acid
12. 4-nitrophenol
13. 4-Chloracetanilide
14. 2-nitrophenol


Mobile phase A:
10mM ammonium formate pH3.0
Mobile phase B:
10mM ammonium formate pH3.0 in ACN
Flow rate : 0.4ml/min
Wavelength : 254nm
Temperature : 40°C

* All columns 3µm 100x2.1mm


Method development screening kits

When developing a new method in chromatography having a diverse range of selectivities allows a choice to be made dependant upon initial knowledge of the compound types and classes: choose phases based on similarity i.e. Evosphere C18/AR and C18/PFP both having a high hydrophobicity, but subtle changes in steric terms. Or choose stationary phases that are as orthogonal as possible from each other allowing for the best probability of a generic gradient screen to ascertain the best starting column to then be taken forward for further optimisation.


**C12
Diphenyl
C18/PFP**



**AQUA
C18/AR
RP18-Amide**



**PFP
Phenyl-Hexyl**




Evosphere C18/AR




- Evosphere C18/AR
- Orthogonal Selectivity
- Method development starting point

Evosphere C18/AR is designed to provide increased resolution between compounds, having a combination of hydrophobicity and aromatic selectivity will lead to enhanced resolution. USP L1 column.




Evosphere Selectivity

When developing a new method in chromatography having a diverse range of selectivities to choose from can help in deciding how peaks resolve and which is the best starting point. In this example Evosphere C18/AR highlights orthogonal selectivity for acid and basic molecules.




Evosphere C18/AR and Evosphere C18/PFP are both USP L1 columns, however they are designed to enhance separation capabilities by offering different mechanisms of interaction to each other, leading to orthogonal separations.



Evosphere C18/PFP

EVO SPHERE





- Evosphere C18/PFP
- Orthogonal Selectivity
- Method development starting point

Evosphere C18/PFP is designed to provide characteristics which will enhance method development. It provides the ability to obtain sharp peak shapes whilst retaining and separating a wide variety of compounds both hydrophobic and hydrophilic. USP L1 column.

Evosphere Selectivity


When developing a new method in chromatography having a diverse range of selectivities to choose from can help in deciding how peaks resolve and which is the best starting point. In this example Evosphere C18/PFP highlights orthogonal selectivity for halogenated positional isomers.





- Predominantly Hydrophobic
- Added Steric and Dipole selectivity
- Ideal for mixtures of neutrals and isomeric species

Untargeted Metabolomics


Metabolomics can offer a challenge in conventional chromatography due to the diverse nature of the analytes. Evosphere C18/PFP was chosen due to its combination of regioisomer and hydrophobic selectivity as well as polar retention capacity.



3 μ m Evosphere C18/PFP 100x2.1mm
p/n : EVO18FP-020503




Evosphere C12




- Evosphere C12
- Ultra High Efficiency
- Method development starting point

Evosphere C12 is designed to provide characteristics which will enhance method development. The dense C12 ligand provides the ability to obtain sharp peak shapes whilst retaining and separating a wide variety of acid, base and neutral compounds with excellent robustness.


Lipidomics Characterisation




Lyso PC 16:0 & 18:0



RT:




PCs with different degrees of Unsaturation



Phosphatidylcholine - Evosphere is able to separate different degrees of unsaturation


16 & 18 Carbon Lyso PC show excellent peak shape and selectivity



- Predominantly Hydrophobic
- High surface protection = higher pH
- Ideal for mixtures of neutrals and acids and bases


Evosphere High pH Stability

Evosphere C12 is designed to offer hydrophobic retention but with an increase of coverage of ligand, this leads to a high retentive, highly stable stationary phase which can be used in new method development. Its selectivity has been used to provide sharper peak shapes due to the high surface protection also adds increased high pH stability to the phase.



5µm Evosphere C12 50x3.0mm Mobile phase: 70:30 0.1% NH₃ : ACN



Evosphere RP18-Amide



- Evosphere RP18-Amide
- Orthogonal Selectivity
- Excellent method development option


Evosphere RP18-Amide is designed to provide polar characteristics which will enhance resolution in method development. It provides orthogonal selectivity to alkyl chain phases due to its polar-embedded group. Sharp peak shapes, extra selectivity and increased retention can all be obtained.

PFAS Selectivity




Excellent Peak Shapes

Evosphere RP18-Amide will offer the sharpest peak shapes for basic analytes. The positive charge of the amide functionality in the stationary phase stopping any tailing from occurring. This stationary phase will also give strong retention of small polar acid* compounds due to their interaction with the positively charged amide functional group (*see apps guide).



Evosphere Diphenyl





- Evosphere Diphenyl
- Separate positional isomers
- Stable ligand, No “MS” bleed

Evosphere Diphenyl is designed to provide pi-pi, steric and hydrophobic characteristics which will enhance selectivity and the ability to develop methods. Particularly suited to positional isomers and other closely related species such as metabolites.


Enhanced Selectivity

When compounds are positional isomers or similar functionality then Evosphere Diphenyl brings extra selectivity over more hydrophobic stationary phases. Having pi-pi interactions as well as a steric selectivity term due to the branched structure allows the resolution of critical pairs.



Isomeric Selectivity

When compounds are positional isomers then Evosphere Diphenyl brings extra selectivity over more hydrophobic stationary phases. Having 3 modes of interaction, pi-pi, steric selectivity and hydrophobicity allows for extra retention and orthogonal selectivity. It is critical that isomers are separated in the LC as MS will struggle to differentiate between them. Fast analysis can still be achieved even with closely related species if sufficient resolution is achieved by the high efficiency monodisperse particles.





Evosphere AQUA



- Evosphere AQUA
- Separate polar species
- Excellent stability


Evosphere AQUA is designed to provide characteristics which will enhance retention of highly polar analytes. Reproducible surface characteristics provide robust separations. A combination of hydrophobic and hydrophilic nature.

Mycotoxins




Nitrosamines

Nitrosamines must be monitored for their presence as they are widely suspected of being carcinogenic. Hundreds of nitrosamines exist and they can vary widely in their chemical nature being hydrophilic or hydrophobic in nature. This presents a challenge when developing a method that can function for many of these groups. Evosphere AQUA having both polar nature and hydrophobic can retain diverse compound sets such as these.



Evosphere Phenyl-Hexyl




- Evosphere Phenyl-Hexyl
- Separate metabolites
- Excellent resolution

Evosphere Phenyl-Hexyl is designed to provide characteristics which will enhance selectivity. It provides alternate selectivity to a pure hydrophobic stationary phase whilst still maintaining the key attributes of robustness and reproducibility.

Enhanced Selectivity

Phenyl-Hexyl offers yet another alternative selectivity in the Evosphere family, combining a short alkyl chain with a phenyl functionality. In this example the resolution of three aromatic nitro compounds is highlighted in a simple mobile phase. The stationary phase providing excellent selectivity of the 3 derivatives.



Evosphere PFP




- Evosphere PFP
- Orthogonal Selectivity
- Combined with Ultra High Efficiency particles

Evosphere PFP (PentaFluoroPhenyl) is designed to provide characteristics which will enhance selectivity. It provides alternate selectivity to a hydrophobic stationary phase whilst still maintaining the key attributes of robustness and reproducibility.

Enhanced Selectivity

Evosphere PFP is complementary to the alkyl-chain aromatic stationary phases since it provides strong electronegative fluorine atoms resulting in strong retention of halogenated and polar species. Evosphere PFP can also aid in the separation of isomeric species due to ability for shape selectivity with its rigid bonded phase.



Evosphere BIO




- Evosphere BIO Monodisperse particles
- 300Å for larger peptide and proteins
- High Efficiency, high sensitivity particle

Evosphere BIO is designed for those compounds that are larger than 2000Da and require a large pore diameter in order to have access to the stationary phase for accurate adsorption/desorption mechanisms.

Evosphere BIO

When analysing complex samples Evosphere minimises band dispersion due to its monodisperse nature, this has been shown to provide peak widths half of other commercial columns, leading to better resolution, better peak height and better sensitivity of low abundance peptides and proteins.

Evosphere C18/AR adds a new dimension to selectivity for these species.




Evosphere BIOMAX

300Å Evosphere in Inert column hardware

Many peptides and proteins do not interact well with traditional LC column hardware since it is stainless steel, generally with stainless steel frits holding the stationary phase in place.

BIOMAX columns are passivated with a new bio-inert coating in order to prevent unnecessary interactions with peptides and proteins, allowing for high sensitivity and full recovery of all analytes. Whilst PEEK can be used, PEEK can swell under pressure so is not ideal in the UHPLC methods we wish to design. BIOMAX removes this issue so that 1.7µm particles can be used for the ultimate in sensitivity and resolution.









Evosphere BIO physicals

	Particle Size	Surface Area	Pore Size	% C	pH range	USP
Evosphere BIO C12	1.7µm 3µm 5µm	n/a	300Å	5%	1-9	L87
Evosphere BIO Diphenyl	1.7µm 3µm 5µm	n/a	300Å	5%	2-9	L11
Evosphere BIO C4	1.7µm 3µm 5µm	n/a	300Å	3%	2-9	L26
Evosphere BIO C18/AR	1.7µm 3µm 5µm	n/a	300Å	7%	2-9	L1

Applications




PFAS - PERFLUOROALKYL SUBSTANCES




Evosphere C18/AR
1.7 μ m 100x2.1mm

Evosphere RP18-Amide
1.7 μ m 100x2.1mm

43 PESTICIDES



CATECHOLAMINES



Column: 3 μ m Evosphere AQUA 150x4.6mm

Mobile Phase: A - 0.1% formic acid

B - 0.1% formic acid in ACN

0-70% B in 10mins

Temp: 20°C

Wavelength : 270nm


1. Dopamine

2. Serotonin

3. DOPAC

4. 5-HIAA

XANTHINE DERIVATIVES



Column: 5 μ m Evosphere C12 150x4.6mm

Mobile Phase: 70:30 0.1% formic acid in Water : MeOH

Flow: 1.0ml/min

Wavelength : 254nm

1. Theobromine

2. Theophylline


3. Caffeine

Applications



OLIGONUCLEOTIDE WITHOUT ION PAIR REAGENT

Separation of Sequence Isomers




- 1 - ATCGATCGATCGATCG**G**
- 2 - ATCGATCGATCGATCG**T**
- 3 - ATCGATCGATCGATCG**C**
- 4 - ATCGATCGATCGATCG**A**

1.7 μ Evosphere MAX C18/AR 100x2.1mm

25mM Ammonium acetate pH=6

12-20% MeOH in 10min

Temp: 60°C



- 1 - ATCGATCGA**A**CGATCGATCG
- 2 - ATCGATCGA**T**AGATCGATCG
- 3 - ATCGATCGATCGATCGA**ACG**
- 4 - ATCGATCGATCGATCG**A**TCA


1.7 μ Evosphere MAX C18/AR 100x2.1mm

25mM Ammonium acetate pH=6

12-20% MeOH in 10min

Temp: 60°C


ISOMERIC PROGESTERONES



Evosphere C18/PFP
5 μ 150x4.6mm
p/n: EVO18FP-050705
Mobile phase:
30:70 A: 0.1% formic acid in water
B: MeOH
Flow: 1.0ml/min
Temp: 20°C
Wavelength: 254nm

- 1. 21-Hydroxyprogesterone
- 2. 17 α -Hydroxyprogesterone

ANTIARRHYTHMIC DRUGS




Column: 5 μ m Evosphere C18/AR 150x4.6mm

Mobile Phase: 70:30 0.1% formic acid in Water : MeOH
Flow: 1.0ml/min
Temp: 25°C
Wavelength : 235nm

- 1. Quinidine
- 2. Hydroquinidine

ISOMERIC ACIDS



Column: 5 μ m Evosphere AQUA 150x4.6mm


Mobile Phase: 0.1% formic acid in Water
Flow: 1.0ml/min
Temp: 20°C
Wavelength : 254nm

- 1. Isoascorbic acid
- 2. Ascorbic acid

Applications



HALOGENATED POSITIONAL ISOMERS



Mobile Phase: 50:50 Water:MeOH

Flow: 1.0ml/min

Temp: 20°C

Wavelength: 254nm


1. Acetophenone

2. 2-Chloroacetophenone

3. 4-Chloroacetophenone

4. 3-Chloroacetophenone

SUBSTITUTED BENZENES



Mobile Phase: 50:50 Water:MeOH

Flow: 1.0ml/min

Temp: 20°C

Wavelength: 210nm

1. 1,2,3-Trimethoxybenzene

2. 1,2-Dimethoxybenzene

3. 1,2,4-Trimethoxybenzene

4. 1,4-Dimethoxybenzene


5. Anisole

6. 1,3-Dimethoxybenzene


7. 1,3,5-Trimethoxybenzene

8. Toluene

SELECTIVITY COMPARISON - EXPLOSIVES



NUCLEOSIDES



Column: 3µm Evosphere AQUA 150x4.6mm

Mobile Phase: 98:2 25mM NH₄OAc : ACN

Flow: 1.0ml/min

Temp: 20°C

Wavelength: 254nm


1. Uracil

2. Uridine

3. Cytosine

4. Guanosine

STEROIDS



Mobile Phase: 30-100% B in 10mins

A: 0.1% Formic acid in Water

B: 0.1% Formic acid in ACN

Flow: 1.0ml/min

Wavelength: 254nm

1. Prednisolone

2. Prednisone


3. Cortisone

4. 17α Hydroxyprogesterone


Applications



XANTHINE DERIVATIVES




ZEARELENONE



Mobile Phase: A: 0.1% formic acid
B: ACN
50-80% B in 5mins, hold for 5
Flow: 1.0ml/min
Temp: 20°C
Wavelength: 250nm

1. Zearalenone


ACIDS



Mobile Phase: 80:20 20mM KH₂PO₄ (pH 2.4) : MeCN
Flow: 1.0ml/min
Temp: 20°C
Wavelength: 254nm

1. Benzoic acid
2. Sorbic acid


TRIFLURALIN



Mobile Phase: 85:15 MeOH: Water
Flow: 1.2ml/min
Temp: 20°C
Wavelength: 254nm

1. Trifluralin

NICOTINAMIDES



Mobile Phase: 98:2 0.1% formic acid in Water : MeOH
Flow: 1.0ml/min
Temp: 20°C
Wavelength: 254nm

1. Isonicotinamide
2. Isonicotinic acid
3. Nicotinamide
4. Nicotinic acid

Applications

Compound	Use	Column		
1,2 Dimethoxybenzene		Evosphere C18/PFP	Amitriptyline Hydrochloride	Antidepressants
1,2,3 Trimethoxybenzene		Evosphere C18/PFP	Ascorbic acid	Vitamins
1,2,4 Dimethoxybenzene		Evosphere C18/PFP	Aspirin	Method Development
1,3 Dinitrobenzene	Explosives	Evosphere Phenyl-Hexyl	Atorvastatin	High Blood Pressure
1,3 Dinitrobenzene	Explosives	Evosphere C18/AR	Bendroflumethiazide	Diuretic
1,3 Dinitrobenzene	Explosives	Evosphere PFP	Benfluralin	Herbicide
1,3,5 Dinitrobenzene	Explosives	Evosphere RP18-Amide	Benzene sulphonic acid	
1,3,5 Dinitrobenzene	Explosives	Evosphere Phenyl-Hexyl	Benzoin	Preservatives in food
1,3,5 Dinitrobenzene	Explosives	Evosphere C18/AR	Benzophenone	Phenones
1,3,5 Dinitrobenzene	Explosives	Evosphere PFP	Benzyl alcohol	
1,4 Dimethoxybenzene		Evosphere C18/PFP	Benzyl alcohol	
17 α -Hydroxyprogesterone		Evosphere AQUA	Benzyl butyl phthalate	Plasticizers
17 α -Hydroxyprogesterone	Hormones	Evosphere C18/PFP	Bis(2-ethylhexyl) adipate	Plasticizers
1-Nitrosopiperidene	Nitrosamines	Evosphere AQUA	Bis(2-ethylhexyl) phthalate	Plasticizers
1-Nitrosopyrrolidine	Nitrosamines	Evosphere AQUA	Butyrophenone	Phenones
21-Hydroxyprogesterone	Hormones	Evosphere C18/PFP	Caffeine	Flu Relief
2-Acetamidophenol	Method Development	Evosphere C18/PFP	Caffeine	Stimulants
2-Aminophenol	Amino Acids	Evosphere C18/AR	Caffeine	Stimulants
2-Aminophenol	Amino Acids	Evosphere C18/AR	Caffeine	Method Development
2-Chloroacetophenone		Evosphere C18/PFP	Cefaclor	
2-Hydroxybenzoic acid	Method Development	Evosphere C18/PFP	Cefadroxil	
2-Hydroxyestradiol	Steroids	Evosphere C18/AR	Cefradine	
2-Nitrophenol	Method Development	Evosphere C18/PFP	Cephalexin	
3,4-Dihydroxyphenyl acetic acid	Amino Acids	Evosphere C18/AR	Chloramphenicol	Eye antibiotic
3-Chloroacetophenone		Evosphere C18/PFP	Cholesterol Esters	Lipidomics
3-methoxycatecol		Evosphere AQUA	Clomipramine Hydrochloride	Antidepressants
3-Methoxy-p-Tyramine	Amino Acids	Evosphere C18/AR	Clomipramine Hydrochloride	Antidepressants
3-methyl-4-nitrobenzoic acid		Evosphere PFP	Co-codamol	Pain relief
3-Methylindole	Amino Acids	Evosphere C18/AR	Cortisone	Steroid
4-Chloracetanilide	Method Development	Evosphere C18/PFP	Cortisone	Steroid
4-Chloroacetophenone		Evosphere C18/PFP	Cytosine	Nucleosides
4-Hydroxybenzoic acid	Method Development	Evosphere C18/PFP	Diazepam	Anti anxiety
4-Hydroxyestradiol	Steroids	Evosphere C18/AR	Diazepam	Anti anxiety
4-methylcatechol		Evosphere AQUA	Dibutyl phthalate	Plasticizers
4-nitrocatechol		Evosphere AQUA	DOPAC	Catecholamines
4-Nitrophenol	Method Development	Evosphere C18/PFP	Dopamine	Catecholamines
5-Aminopentanoic acid	Metabolomics	Evosphere C18/PFP	Dopamine	Amino Acids
5-HIAA	Catecholamines	Evosphere AQUA	Doxepin	Antidepressants
5-Hydroxy Tryptophan	Amino Acids	Evosphere C18/AR	Folic Acid	Vitamin
5-Methoxytryptaphol	Amino Acids	Evosphere C18/AR	Furazolidone	Antibacterials
Acetaminophen		Evosphere PFP	GABA	Amino Acids
Acetaminophen	Flu Relief	Evosphere AQUA	Guanosine	Nucleosides
Acetaminophen	Pain relief	Evosphere AQUA	Hydroquinidine	Irregular heartbeat treatment
Acetophenone	Phenones	Evosphere AQUA	Hydroquinidine	Irregular heartbeat treatment
Acetophenone		Evosphere C18/PFP	Hydroquinidine	Method Development
Acetyl choline	Amino Acids	Evosphere C18/AR	Ibuprofen	Anti-inflammatory painkillers
Acetyl-Carnitine	Metabolomics	Evosphere C18/PFP	Ibuprofen	Anti-inflammatory painkillers
Aflatoxin B1	Mycotoxins	Evosphere AQUA	Ibuprofen	Anti-inflammatory painkillers
Aflatoxin B2	Mycotoxins	Evosphere AQUA	Imipramine	Antidepressants
Aflatoxin G1	Mycotoxins	Evosphere AQUA	Imipramine Hydrochloride	Antidepressants
Aflatoxin G2	Mycotoxins	Evosphere AQUA	Imipramine Hydrochloride	Antidepressants
Amitriptyline Hydrochloride	Antidepressants	Evosphere Phenyl-Hexyl	Indole-3-Carboxylic Acid	Amino Acids
			Indomethacin	Anti-inflammatory painkillers
			Indomethacin	Anti-inflammatory painkillers
			Indomethacin	Anti-inflammatory painkillers
			Inole-3-Acetamide	Amino Acids
			Iso-ascorbic acid	Vitamins
			Isoleucine	Metabolomics
			Iso-leucine	Amino Acids
			Isonicotinamide	Vitamins
			Isonicotinic acid	Vitamins
			Kynurenic Acid	Amino Acids
			Leucine	Metabolomics
			Lorazepam	Anti anxiety
			Lorazepam	Anti anxiety
			L-Phenylalanine	Amino Acids
			Lyso-PC	Lipidomics
			Melatonin	Amino Acids
			Methionine	Amino Acids
			Mianserin Hydrochloride	Antidepressants
			Mianserin Hydrochloride	Antidepressants
			N-acetyl Tyrosine EE	Amino Acids
			N-Acetyl-4-Hydroxytryptamine	Amino Acids
			N-acetyl Tryptophan EE	Amino Acids
			Nalidixic acid	Antibacterials
			Nicotinamide	Vitamins
			Nicotinic acid	Vitamins
			Nicotinic Acid	Amino Acids
			Nitrobenzene	Explosives
			Nitrobenzene	Explosives
			N-Nitrosodibutylamine	Nitrosamines
			N-Nitrosodiethylamine	Nitrosamines
			N-Nitrosodimethylamine	Nitrosamines
			N-Nitrosodi-n-propylamine	Nitrosamines
			N-Nitrosomethylamine	Nitrosamines
			Norepinephrine	Amino Acids
			Nortriptyline	Antidepressants
			Oligonucleotide	Oligonucleotide
			Paracetamol	Method Development
			Patulin	Mycotoxin in molds
			PFHPS	PFAS
			PFHPS	PFAS
			PFHXS	PFAS
			PFHXS	PFAS
			PFOA	PFAS
			PFOA	PFAS
			PFOS	PFAS
			PFOS	PFAS
			Phenol	
			Phenol	Method Development
			Phenylalanine	Metabolomics
			Phenylephrine	Flu Relief
			Phospholipids	Lipidomics
			Prednisolone	
			Prednisolone	
			Prednisone	
			Prednisone	
			Propiophenone	Phenones
			Quinidine	Irregular heartbeat treatment
			Quinidine	Irregular heartbeat treatment
			Quinolinic Acid	Amino Acids
			Quinolinic Acid	Amino Acids
			Resorcinol	
			Serotonin	Catecholamines
			Serotonin	Amino Acids
			Sorbitic acid	Preservatives in food
			Sulfamerazine	
			Sulfamerazine	
			Sulfamethoxazole	
			Sulfamethoxazole	
			Sulfathiazole	
			Sulfathiazole	
			Temazepam	Anti anxiety
			Temazepam	Anti anxiety
			Theobromine	Stimulants
			Theobromine	Stimulants
			Theobromine	Method Development
			Theophylline	Stimulants
			Theophylline	Stimulants
			Theophylline	Method Development
			Trifluralin	Herbicide
			Trifluralin	Herbicide
			Triglycerides	Lipidomics
			Trimipramine	Antidepressants
			Tryptanthrin	Amino Acids
			Tryptophan	Amino Acids
			Tryptophan Ethyl Ester	Amino Acids
			Tryptophan ME	Amino Acids
			Tryptophan Methyl Ester	Amino Acids
			Tyramine	Amino Acids
			Tyrosine	Metabolomics
			Uracil	Nucleosides
			Uridine	Nucleosides
			Valine	Metabolomics
			Zearalenone	Mycotoxin in molds
			Zearalenone	Mycotoxin in molds

Applications

Capillaries & Scaling to Prep



Evosphere capillaries are available in 75 μ m, 150 μ m, 0.5mm, 1mm i.d. with any phase chemistry and any particle size from the Evosphere range. Request a quote from your local distributor.



Evosphere Prep

- 10mm, 21.2mm and 30mm
- High Loadability
- Optimised Packing Efficiency
- Narrow peak profile, High Efficiency and Resolution

Evosphere Prep columns are designed for high sample loading, high throughput applications. The optimised packed bed (OPB) process ensures excellent peak shapes and efficiency, whilst the lifetime of the column is increased.





MAX Hardware

Inert Column Hardware Design

Many peptides and proteins do not interact well with traditional LC column hardware since it is stainless steel, generally with stainless steel frits holding the stationary phase in place.

MAX columns are passivated with a new bio-inert coating in order to prevent unnecessary interactions with peptides and proteins, allowing for high sensitivity and full recovery of all analytes. Whilst PEEK can be used, PEEK can also swell under pressure so is not ideal in the UHPLC methods we wish to design. MAX removes this issue so that small 1.7 μ m particles can be used for the ultimate in sensitivity and resolution.



1.7µm EVOSPHERE® part numbers

1.7µm EVOSPHERE C12		Column Length			
		30	50	100	150
	2.1	EV012-020201	EV012-020301	EV012-020501	EV012-020701
Column Diameter	3.0	EV012-030201	EV012-030301	EV012-030501	EV012-030701
	4.6	EV012-050201	EV012-050301	EV012-050501	EV012-050701

1.7µm EVOSPHERE C18/AR		Column Length			
		30	50	100	150
	2.1	EV018AR-020201	EV018AR-020301	EV018AR-020501	EV018AR-020701
Column Diameter	3.0	EV018AR-030201	EV018AR-030301	EV018AR-030501	EV018AR-030701
	4.6	EV018AR-050201	EV018AR-050301	EV018AR-050501	EV018AR-050701

1.7µm EVOSPHERE C18/PFP		Column Length			
		30	50	100	150
	2.1	EV018FP-020201	EV018FP-020301	EV018FP-020501	EV018FP-020701
Column Diameter	3.0	EV018FP-030201	EV018FP-030301	EV018FP-030501	EV018FP-030701
	4.6	EV018FP-050201	EV018FP-050301	EV018FP-050501	EV018FP-050701

1.7µm EVOSPHERE RP18-AMIDE		Column Length			
		30	50	100	150
	2.1	EVORP18-020201	EVORP18-020301	EVORP18-020501	EVORP18-020701
Column Diameter	3.0	EVORP18-030201	EVORP18-030301	EVORP18-030501	EVORP18-030701
	4.6	EVORP18-050201	EVORP18-050301	EVORP18-050501	EVORP18-050701

1.7µm EVOSPHERE PHENYL-HEXYL		Column Length			
		30	50	100	150
	2.1	EVOHEX-020201	EVOHEX-020301	EVOHEX-020501	EVOHEX-020701
Column Diameter	3.0	EVOHEX-030201	EVOHEX-030301	EVOHEX-030501	EVOHEX-030701
	4.6	EVOHEX-050201	EVOHEX-050301	EVOHEX-050501	EVOHEX-050701

1.7µm EVOSPHERE DIPHENYL		Column Length			
		30	50	100	150
	2.1	EVOPH-020201	EVOPH-020301	EVOPH-020501	EVOPH-020701
Column Diameter	3.0	EVOPH-030201	EVOPH-030301	EVOPH-030501	EVOPH-030701
	4.6	EVOPH-050201	EVOPH-050301	EVOPH-050501	EVOPH-050701

1.7µm EVOSPHERE PFP		Column Length			
		30	50	100	150
	2.1	EVOPFP-020201	EVOPFP-020301	EVOPFP-020501	EVOPFP-020701
Column Diameter	3.0	EVOPFP-030201	EVOPFP-030301	EVOPFP-030501	EVOPFP-030701
	4.6	EVOPFP-050201	EVOPFP-050301	EVOPFP-050501	EVOPFP-050701

1.7µm EVOSPHERE AQUA		Column Length			
		30	50	100	150
	2.1	EVOAQ-020201	EVOAQ-020301	EVOAQ-020501	EVOAQ-020701
Column Diameter	3.0	EVOAQ-030201	EVOAQ-030301	EVOAQ-030501	EVOAQ-030701
	4.6	EVOAQ-050201	EVOAQ-050301	EVOAQ-050501	EVOAQ-050701

3µm EVOSPHERE® part numbers

3µm EVOSPHERE C12		Column Length			
		30	50	100	150
	2.1	EV012-020203	EV012-020303	EV012-020503	EV012-020703
Column Diameter	3.0	EV012-030203	EV012-030303	EV012-030503	EV012-030703
	4.6	EV012-050203	EV012-050303	EV012-050503	EV012-050703

3µm EVOSPHERE C18/AR		Column Length			
		30	50	100	150
	2.1	EV018AR-020203	EV018AR-020303	EV018AR-020503	EV018AR-020703
Column Diameter	3.0	EV018AR-030203	EV018AR-030303	EV018AR-030503	EV018AR-030703
	4.6	EV018AR-050203	EV018AR-050303	EV018AR-050503	EV018AR-050703

3µm EVOSPHERE C18/PFP		Column Length			
		30	50	100	150
	2.1	EV018FP-020203	EV018FP-020303	EV018FP-020503	EV018FP-020703
Column Diameter	3.0	EV018FP-030203	EV018FP-030303	EV018FP-030503	EV018FP-030703
	4.6	EV018FP-050203	EV018FP-050303	EV018FP-050503	EV018FP-050703

3µm EVOSPHERE RP18-AMIDE	
--------------------------	--

5µm EVOSPHERE® part numbers

5µm EVOSPHERE C12		Column Length			
		30	50	100	150
	2.1	EV012-020205	EV012-020305	EV012-020505	EV012-020705
Column Diameter	3.0	EV012-030205	EV012-030305	EV012-030505	EV012-030705
	4.6	EV012-050205	EV012-050305	EV012-050505	EV012-050705

5µm EVOSPHERE C18/AR		Column Length			
		30	50	100	150
	2.1	EV018AR-020205	EV018AR-020305	EV018AR-020505	EV018AR-020705
Column Diameter	3.0	EV018AR-030205	EV018AR-030305	EV018AR-030505	EV018AR-030705
	4.6	EV018AR-050205	EV018AR-050305	EV018AR-050505	EV018AR-050705

5µm EVOSPHERE C18/PFP		Column Length			
		30	50	100	150
	2.1	EV018FP-020205	EV018FP-020305	EV018FP-020505	EV018FP-020705
Column Diameter	3.0	EV018FP-030205	EV018FP-030305	EV018FP-030505	EV018FP-030705
	4.6	EV018FP-050205	EV018FP-050305	EV018FP-050505	EV018FP-050705

5µm EVOSPHERE RP18-AMIDE		Column Length			
		30	50	100	150
	2.1	EVORP18-020205	EVORP18-020305	EVORP18-020505	EVORP18-020705
Column Diameter	3.0	EVORP18-030205	EVORP18-030305	EVORP18-030505	EVORP18-030705
	4.6	EVORP18-050205	EVORP18-050305	EVORP18-050505	EVORP18-050705

5µm EVOSPHERE PHENYL-HEXYL		Column Length			
		30	50	100	150
	2.1	EVOHEX-020205	EVOHEX-020305	EVOHEX-020505	EVOHEX-020705
Column Diameter	3.0	EVOHEX-030205	EVOHEX-030305	EVOHEX-030505	EVOHEX-030705
	4.6	EVOHEX-050205	EVOHEX-050305	EVOHEX-050505	EVOHEX-050705

5µm EVOSPHERE DIPHENYL		Column Length			
		30	50	100	150
	2.1	EVOPH-020205	EVOPH-020305	EVOPH-020505	EVOPH-020705
Column Diameter	3.0	EVOPH-030205	EVOPH-030305	EVOPH-030505	EVOPH-030705
	4.6	EVOPH-050205	EVOPH-050305	EVOPH-050505	EVOPH-050705

5µm EVOSPHERE PFP		Column Length			
		30	50	100	150
	2.1	EVOPFP-020205	EVOPFP-020305	EVOPFP-020505	EVOPFP-020705
Column Diameter	3.0	EVOPFP-030205	EVOPFP-030305	EVOPFP-030505	EVOPFP-030705
	4.6	EVOPFP-050205	EVOPFP-050305	EVOPFP-050505	EVOPFP-050705

5µm EVOSPHERE AQUA		Column Length			
		30	50	100	150
	2.1	EVOAQ-020205	EVOAQ-020305	EVOAQ-020505	EVOAQ-020705
Column Diameter	3.0	EVOAQ-030205	EVOAQ-030305	EVOAQ-030505	EVOAQ-030705
	4.6	EVOAQ-050205	EVOAQ-050305	EVOAQ-050505	EVOAQ-050705

- Direct connect guard system for all 3µm and 5µm phases
- Quick replacement cartridges
- Highly Cost Effective



5µm Evosphere Guard Cartridges	
DCGUA-1	Guard Cartridge Holder
DCxx-040005G/2	10x4mm Evosphere 5µm Guard pk 2
DCxx-040005G/4	10x4mm Evosphere 5µm Guard pk 4
DCxx-020005G/2	10x2mm Evosphere 5µm Guard pk 2
DCxx-020005G/4	10x2mm Evosphere 5µm Guard pk 4

Replace xx 12 for Evosphere C12 EPH for Evosphere Diphenyl AQ for Evosphere AQUA HEX for Evosphere Phenyl-Hexyl
PFP for Evosphere PFP RP18 for Evosphere RP18-Amide 18AR for Evosphere C18/AR 18FP for Evosphere C18/PFP

EVOSPHERE® BIO part numbers





EVOSPHERE BIO C12		Column Length			
		30	50	100	150
	2.1	EVOBIO312-0202xx	EVOBIO312-0203xx	EVOBIO312-0205xx	EVOBIO312-0207xx
Column Diameter	3.0	EVOBIO312-0302xx	EVOBIO312-0303xx	EVOBIO312-0305xx	EVOBIO312-0307xx
	4.6	EVOBIO312-0502xx	EVOBIO312-0503xx	EVOBIO312-0505xx	EVOBIO312-0507xx

EVOSPHERE BIO DIPHENYL		Column Length			
		30	50	100	150
	2.1	EVOBIO3PH-0202xx	EVOBIO3PH-0203xx	EVOBIO3PH-0205xx	EVOBIO3PH-0207xx
Column Diameter	3.0	EVOBIO3PH-0302xx	EVOBIO3PH-0303xx	EVOBIO3PH-0305xx	EVOBIO3PH-0307xx
	4.6	EVOBIO3PH-0502xx	EVOBIO3PH-0503xx	EVOBIO3PH-0505xx	EVOBIO3PH-0507xx

EVOSPHERE BIO C4		Column Length
------------------	--	---------------



Other Product Guides Available



WORLDWIDE AVAILABILITY



45 Coalbrookdale Road
Clayhill Industrial Park
Neston
Cheshire, UK
CH64 3UG

t: +44 151 336 2266
f: +44 151 336 2669
www.fortis-technologies.com
e: info@fortis-technologies.com

Company No. 5449466
VAT No. 866 8966 43

Fortis products are available worldwide. For the distributor in your country, contact Fortis international Sales Office, UK by telephone, fax or email: info@fortis-technologies.com

- Austria
- Hong Kong
- Poland
- Bangladesh
- Hungary
- Portugal
- Brazil
- India
- Romania
- Canada
- Ireland
- Russia
- China
- Israel
- Singapore
- Columbia
- Italy
- South Africa
- Czech Republic
- Japan
- Spain
- Ecuador
- Korea
- Sweden
- Egypt
- Malaysia
- Switzerland
- France
- Mexico
- Taiwan
- Germany
- Netherlands
- Thailand
- Greece
- Norway
- Turkey
- Holland
- Puerto Rico
- USA

For technical support or applications contact :
technicalsupport@fortis-technologies.com

For more information VISIT :
www.fortis-technologies.com

Sign up to our Newsletter here :



Join our Linked-In page here :

